

This is a repository copy of *Aragonite bias exhibits systematic spatial variation in the late Cretaceous Western Interior Seaway, North America*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/150559/>

Version: Accepted Version

Article:

Dean, Christopher, Allison, Peter A., Hampson, Gary et al. (1 more author) (2019) Aragonite bias exhibits systematic spatial variation in the late Cretaceous Western Interior Seaway, North America. *Paleobiology*. ISSN 1938-5331

<https://doi.org/10.1017/pab.2019.33>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Paleobiology



CAMBRIDGE
UNIVERSITY PRESS

Aragonite bias exhibits systematic spatial variation in the late Cretaceous Western Interior Seaway, North America

Journal:	<i>Paleobiology</i>
Manuscript ID	PAB-OR-2019-0021.R2
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Dean, Christopher; Imperial College London, Earth Science and Engineering Allison, Peter; Imperial College London, Earth Science and Engineering Hampson, Gary; Imperial College London, Earth Science and Engineering Hill, Jon; University of York, Department of Environment and Geography
Geographic Location:	United States
Taxonomy:	Mollusca, Bivalvia, Cephalopoda
Analysis:	diversity pattern, geographic distribution, multiple regression
Geologic Age:	Campanian, Cenomanian, Turonian
Topic:	bias, taphonomy, biodiversity

SCHOLARONE™
Manuscripts

Aragonite bias exhibits systematic spatial variation in the late Cretaceous Western Interior Seaway, North America.

Christopher D. Dean¹, Peter A. Allison¹, Gary J. Hampson¹ and Jon Hill²

¹Department of Earth Science and Engineering, Imperial College London, London, United Kingdom;

²Department of Environment and Geography, University of York, Heslington, York, United Kingdom;

e-mail: christopher.dean09@imperial.ac.uk

ABSTRACT

Preferential dissolution of the biogenic carbonate polymorph aragonite promotes preservational bias in shelly marine faunas. Whilst field studies have documented the impact of preferential aragonite dissolution on fossil molluscan diversity, its impact on regional and global biodiversity metrics is debated. Epicontinental seas are especially prone to conditions which both promote and inhibit preferential dissolution, which may result in spatially extensive zones with variable preservation. Here we present a multi-faceted evaluation of aragonite dissolution within the late Cretaceous Western Interior Seaway of North America. Occurrence data of molluscs from two time intervals (Cenomanian-Turonian boundary, early Campanian) are plotted on new high-resolution paleogeographies to assess aragonite preservation within the seaway. Fossil occurrences, diversity estimates and sampling probabilities for calcitic and aragonitic fauna were compared in zones defined by depth and distance from the seaway margins. Apparent range sizes, which could be influenced by differential preservation potential of aragonite between separate localities, were also compared. Our results are consistent with exacerbated aragonite dissolution within specific depth zones for both time slices, with aragonitic bivalves additionally showing a statistically significant decrease in range size compared to calcitic fauna within carbonate-dominated Cenomanian-Turonian strata. However, we are unable to conclusively show that aragonite dissolution impacted diversity estimates. Therefore, whilst aragonite dissolution is likely to have affected the preservation of

fauna in specific localities, time averaging and instantaneous preservation events preserve regional biodiversity. Our results suggest that the spatial expression of taphonomic biases should be an important consideration for paleontologists working on paleobiogeographic problems.

Key words: Mollusca, calcite, OAE2, Cretaceous, fossil record bias, sampling bias.

INTRODUCTION

WHILST the fossil record provides a unique window into past life on Earth, it is well known that it is both pervasively and non-uniformly biased (Raup, 1976; Koch, 1978; Foote and Sepkoski, 1999; Alroy et al., 2001; Allison and Bottjer, 2011). Geologic, taphonomic and anthropogenic biases (such as the amount of available fossiliferous rock for sampling, variation in fossilization, and the degree to which the available rock record has been sampled) skew or remove information from the fossil record, leaving the remaining catalogue of data uneven and incomplete. Although biomineralized remains have an increased preservation potential compared to soft bodied tissues (Allison, 1988; Briggs, 2003), they are still influenced by various geologic and taphonomic processes (Kidwell and Bosence, 1991; Kidwell and Brenchley, 1994; Kidwell and Jablonski, 1983; Best, 2008; Hendy, 2011). Shelly marine faunas are especially susceptible to misrepresentation due to preferential dissolution of biogenic carbonate polymorphs. It is well established that aragonite, a polymorph of CaCO_3 found within the biomineralized shells of many invertebrates, dissolves more rapidly than the more common form of CaCO_3 , calcite, and at a higher pH (Canfield and Raiswell, 1991; Tynan and Opdyke, 2011). Whilst both polymorphs can be destroyed by adverse conditions near the sediment-water interface (Best and Kidwell, 2000) and the effects of dissolution can vary between fauna (due to microstructure surface area, morphology, and shell organic content:

Walter and Morse, 1984; Harper, 2000; Kosnik et al., 2011), it is still the case that aragonitic shells are more likely to dissolved than calcitic remains (Brett and Baird, 1986). As mineral composition of molluscs is usually conserved at the Family level (Carter, 1990), this has the potential to skew the record of molluscan diversity and trophic structure through time (Cherns et al., 2011; Cherns and Wright, 2000; Wright et al., 2003), and negatively affect subsequent work that relies on the relative abundance and distribution of shelly marine fauna (Kidwell, 2005). Cherns and Wright (2001) argued that early-stage dissolution could be substantial and referred to the phenomenon as the “Missing Mollusc” bias. Subsequent work on a multitude of temporal and spatial scales (Wright et al., 2003; Bush and Bambach, 2004; Kidwell, 2005; Crampton et al., 2006; Valentine et al., 2006; Foote et al., 2015; Jordan et al., 2015, Hsieh et al., 2019) has debated the magnitude of this bias; however, there is a broad agreement on the potential for dominantly aragonitic shells to suffer greater post-mortem diagenetic destruction in the Taphonomically Active Zone (TAZ) (Davies et al., 1989; Foote et al., 2015). Whilst the effects of dissolution on the global macroevolutionary record of molluscs has been found to be limited, possibly due to the potential of aragonite to recrystallize to calcite (Kidwell 2005; Paul et al., 2008; Jordan et al., 2015), it is conceivable that local or regional conditions could impact severely on perceived patterns of biodiversity in restricted areas (Bush and Bambach 2004). In a regional study of Cenozoic molluscs, Foote et al. (2015) found evidence to suggest that aragonite dissolution was both enhanced in carbonate sediments and insignificant within siliciclastic sediments, with similar preservation potential of aragonitic and calcitic fauna within the latter. They further emphasized the fact that scale is an important factor in determining the observable impacts of aragonite dissolution, which will strongly vary between local (potentially consisting of an individual bed), regional and global studies. However, to date research has focused on assessing the influence of aragonite bias on temporal trends of biodiversity and has ignored the potential for direct spatial expression.

77 Early stage dissolution occurs within modern environments as a result of microbially mediated
78 reactions increasing local acidity (Walter et al., 1993; Ku et al., 1999; Sanders, 2003).
79 Bacterially-mediated decay of organic material within the upper sedimentary column occurs in
80 a series of preferential redox reactions. By-products of these reactions, such as solid phase
81 sulphides from sulphate reduction and CO₂ from aerobic oxidation, result in changes to local
82 pore-water saturation of calcium carbonate (Canfield and Raiswell, 1991; Ku et al., 1999).
83 Additionally, oxidation of H₂S above the oxycline increases acidity at that boundary; if this
84 occurs at the sediment-water interface then it can adversely affect the preservation of shelly
85 marine fauna (Ku et al., 1999). As such, dysoxic sedimentary environments might have a
86 predisposition for dissolution of biogenic carbonate and enhance the effect of the “Missing
87 Mollusc” bias (Jordan et al., 2015). Epicontinental seas, marine water-bodies which form by
88 the flooding of continental interiors, are especially prone to strong water column stratification
89 and sea level variation, and have a pre-disposition to seasonally anoxic or dysoxic conditions
90 (Allison and Wells, 2006; Peters, 2009). As such, they have the potential to be more prone to
91 both preferential aragonite loss and preservation than modern oceans. Cherns et al.'s (2011)
92 model for taphonomic gradients of aragonite preservation along a shelf to basin transect can be
93 readily applied to epicontinental sea settings (Fig. 1). If we assume the center of a seaway was
94 stratified with at least a seasonally anoxic basin floor, we would expect enhanced dissolution
95 to occur in the seaway margins, likely in the mid-to-outer shelf setting (Cherns et al., 2011). In
96 the anoxic basin center we would expect to see enhanced preservation, as an aragonitic skeleton
97 residing on the surface sediment in an anoxic water column would not be susceptible to
98 dissolution from H₂S oxidation (Jordan et al., 2015; however, we would not expect to see
99 abundant benthos in such a setting because of bottom water toxicity). It is apparent this could
100 result in spatially expansive zones with conditions predisposed for heightened aragonite
101 dissolution and preservation (Fig. 1; it is important to note that we do not expect all aragonitic

fauna to be missing from any region of the seaway – merely that a lower relative proportion of aragonitic molluscs be found, due to a reduced probability of an individual site recording their occurrence). How these hypothesized basin-margin to basin-center zones could influence long term patterns of mollusc distribution, preservation and recovery remains to be examined. As epicontinental seas contain the majority of our Phanerozoic fossil record (Allison and Wells, 2006), it is imperative that we understand systematic biases that may specifically affect these settings.

Here we present a spatial investigation of aragonite dissolution within the late Cretaceous Western Interior Seaway (WIS) of North America, using sampling probability estimates and multiple logistic regression to evaluate patterns of spatial distribution in preserved calcitic and aragonitic fauna. We address two key questions: (1) does aragonite bias exhibit systematic spatial variation across the seaway and (2) if so, does this influence perceived patterns of diversity?

MATERIALS AND METHODS

Time Intervals and Paleogeography

The two stratigraphic intervals or time slices (Cenomanian-Turonian and early Campanian) were selected: (1) because of purported dysoxic conditions within their duration; and (2) due to their differences in environment, oceanography and preserved lithology, allowing for comparison of taphonomic regimes. The first interval covers the Cenomanian–Turonian boundary, spanning from the *Dunveganoceras pondi* to *Collignonicerias woollgari* ammonite zone (~94.7–93 Ma) (Cobban et al., 2006). The second interval spans the early Campanian, from the *Scaphites leei* III to *Baculites obtusus* ammonite zones (~83.5–80.58 Ma) (Cobban et

al., 2006). The geologic context of stratigraphic intervals is detailed in Supplementary Information 1.

A global atlas of 1:20,000,000 scale paleogeographic maps, compiled by GETECH plc, formed the basis for new regional-scale, high resolution interpretations for the selected time intervals. The original paleogeographic maps (Markwick, 2007) are underpinned by the GETECH plate model (v1), which is outlined further in Supplementary Information 1. High resolution mapping involved synthesis of stratigraphic, sedimentologic and paleontologic information to produce 1:5,000,000 scale paleogeographies with suggested paleobathymetry. A full list of decisions on paleogeographic reconstructions and key references for each time interval are provided in Supplementary Information 1.

Landward-to-basinward arrangements of *a priori* binned zones for each time-slice were based on average paleobathymetry (Fig. 2). Bathymetric reconstructions were divided into four bins, each of which covers a specific interpreted depth range: Nearshore (<50 m), Proximal Offshore (50-100 m), Distal Offshore (100-150 m) and Basin Center (>150 m). These designations were based on the previously constructed paleobathymetry for the WIS produced by Sageman and Arthur (1994), but match the paleobathymetry in our new maps and represent a reasonably high resolution without being compromised by large changes in shoreline position within our chosen time slices.

Distance-from-paleoshoreline zones (Fig. S1) were constructed based on 50 km intervals from the time-averaged paleo-shoreline position until reaching the basin center, with number of occurrences, collections and total outcrop area plotted per zone. These were generated by constructing a fishnet of points in ArcGIS (ESRI, 2010) using the “Fishnet tool”, which were selected by the “Select By Location” tool with increasing distance in 50 km intervals from the paleoshoreline: the position of the most basinward selected points was used for the bin

boundary. Results for depth zones are used in the main body of this manuscript; distance-from-paleoshoreline zones are available in Supplementary Information 1 and Figures S2, S4, S6 and S7.

Fossil Dataset

A presence-only fossil occurrence dataset of bivalve and ammonite taxa was produced for the selected stratigraphic intervals, collated from personally provided digitized collections from the United States Geological Survey (USGS) and Smithsonian Museum of Natural History (NMNH), as well as downloads from the Paleobiology Database (PBDB; <http://paleobiodb.org>), and iDigBio (<http://www.idigbio.org>). Each occurrence includes taxonomic and geographic locality data, an associated collection with lithologic and geologic information, and modern latitudinal and longitudinal co-ordinates. Data were extensively screened for problematic records and to ensure taxonomic validation (see Supplementary Information 2 for the latter).

The resultant Cenomanian–Turonian dataset contains 5867 occurrences from 2409 localities, with 207 genera, 1549 species, and 3886 specimens identifiable to species level. The early Campanian dataset comprises 2544 occurrences from 1186 localities, recording 163 genera, 1405 species, and 1405 specimens identifiable to species level. Generic level taxonomic diversity was used for all tests; species level results can be found in Supplementary Information 1 and in Supplementary figures S3-6. Full information regarding downloads, sources and screening of data can be found in Supplementary Information 1, and the full dataset found in Supplementary Information 2.

Mineralogy

Bivalve shells are a composite of layered mineral crystallites, which are sheathed by a refractory organic matrix of fibrous protein (Taylor, 1969). As these mineral layers can be comprised of both calcite and aragonite, variation in overall mineral composition must be taken into account when assigning a predominant mineralogy to a specific bivalve Genera. Different scoring mechanisms have been adopted by previous workers to address this issue. Kidwell (2005) used a five-point decimal scoring system from entirely aragonitic (1) to entirely calcitic (3), with three permutations of mineralogy between. Crampton et al. (2006) adopted a simple and effective system of counting organisms as calcitic if they contained a calcitic element that would allow them to be identified to species level. We utilise a combination of these approaches - organisms were scored using the system of Kidwell (2005) to maintain the maximum amount of data, but simplified into binary categories afterwards based on whether they contained sufficient calcitic parts to enhance preservation potential. Note that we have not included either the inner myostracal layer or periostracum in our assignments of mineralogy.

Information on shell composition was predominantly gathered from a personally provided dataset from S. Kidwell (Kidwell, 2005), as well as further studies from Taylor (Taylor, 1969; Taylor and Layman, 1972), Majewske (1974), Carter (1990), Schneider and Carter (2001), Lockwood (2003), Hollis (2008) and Ros-Franch (2009), as well as many papers focussed on single genera or families. For genera for which information regarding shell mineralogy was not available, composition was assigned based on the dominant mineralogy of the family, as composition is highly conservative both amongst species within a genera and amongst genera within a family (Taylor, 1969). In total, 124 bivalve genera were assigned a mineralogy, of which 41 (33%) were achieved using familial relation (Supplementary Information 2).

Life habits

Life habits of bivalves were assembled to allow additional interrogation and interpretation of environmental and sampling regimes. Life habits were separated into the following categories: relation to substrate, mobility and diet. Data for each genera of bivalve were primarily gathered from the NMiTA Molluscan Life Habits Database (Todd, 2017) and the PBDB, with further data collected from the wider literature (Supplementary Information 2).

Outcrop Area

Relevant rock outcrop area was plotted per zone to evaluate broader scale bias influencing patterns of fossil distribution. Outcrop areas for the selected time slices were generated by combining state-wide digitized geologic maps from publicly available USGS downloads and selecting shape files which matched formations found within those time slices. Some State Surveys grouped relevant formations with other partially contemporaneous formations that spanned multiple stages: we chose to include these designations in order to present the maximum possible sampling extent in terms of outcrop area. Outcrop was projected in ArcGIS (ESRI, 2010) using the USA Contiguous Albers Equal Area conic projection, to minimize distortion of distances. Outcrop areas per zone were created by using the “Intersect tool” in the Geoprocessing toolbar in ArcGIS, and area (km²) calculated using the Calculate Geometry function in the attribute table. Outcrop was split into depth zones by using the Intersect tool in ArcGIS (ESRI, 2010). Outcrop area for each zone was calculated by summing the total area of all outcrop polygons within that zone. Collections per zone were counted by exporting occurrences selected in zones in the seaway as shapefiles, then using the “arccgisbinding” package to view and organise the data in R version 3.0.2 (Team, 2017).

Dominant lithology

Each collection was assigned a dominant lithology to allow for comparative testing. If these data were not available, a lithology was assigned from the dominant lithology of the formation, with reference to USGS formation records. Collections were assigned one of the following lithologies (primarily based off original USGS records): siliciclastic mudstone, siliciclastic siltstone, siliciclastic sandstone, conglomerate, ironstone, calcareous mudstone and siltstone, marl, calcarenite, limestone and chalk.

Range Size

If the presence of preferentially destructive zones is affecting the spatial distribution of aragonitic fauna, we might expect to see overall smaller range sizes for aragonitic organisms compared to calcitic organisms (Fig. 3). As such, range size estimates were produced for calcitic and aragonitic bivalves and compared to test if aragonite bias influenced perceived range of aragonitic organisms. Note that ammonites were excluded from this test due to the difference in life habit between them and bivalve fauna: ammonites have a pelagic to nektonic-benthic mode of life (Ritterbush et al., 2014), whilst bivalves are predominantly epifaunal and infaunal.

Geographic locality data for the selected fauna was visualized in ArcGIS (ESRI, 2010). Faunal occurrences were paleo-rotated using the Gtech Plate Model to match the paleogeography of the appropriate stages of the Late Cretaceous. This ensures that tectonic expansion and contraction of the North American plate from the Mesozoic to Recent has a negligible effect on propagating estimation error in range-size reconstructions. Fossil occurrences were projected into ArcGIS using the USA Contiguous Albers Equal Area conic projection. A 10 km buffer was additionally applied to each occurrence point in order to control for any error in paleogeographic or present position of fauna. ArcGIS (ESRI, 2010) was then used to construct convex hull polygons for each taxon, and the spatial analyst tools from this

software calculated the area of each reconstructed polygon. We did not account for landforms within the ranges of any organisms, and thus ignored their area when calculating overall area of ranges. Several vertices for range size polygons appeared on what is classified as land within our paleogeographies; due to rapid changes in shoreline position within the WIS, we decided to keep using these fauna for range size estimations. Myers and Lieberman (2011) showed that relative range sizes for vertebrates in the WIS were not overly affected by resampling occurrence points – consequently, we have not carried out a similar test for this study. Comparisons between the ranges of aragonitic and calcitic fauna were carried out using the Wilcoxon-Mann-Whitney test with continuity correction (Brown and Rothery, 1993). Geographic range data for all applicable taxa are provided in Supplementary Information 2.

Sampling Probability and Multiple Logistic Regression

To be able to further observe differences between aragonitic and calcitic organisms throughout the seaway, we employed a modified version of the sampling probability method used by Foote et al. (2015) (after Foote and Raup, 1996). In this method, the sampling probability of a time bin was generated by compiling a list of all fauna with originations older than that bin and extinctions younger, and then dividing the total number of species found within the bin by that figure. This allows for a sampling probability to be estimated on a per bin, per group basis. Here we devised three variants on this method for application in the spatial realm. It should be made clear that the modified methods utilized in this work come with the caveat that in the spatial realm it is impossible to know if a species was present in a precise location in the past: for instance, if zones A, B, and C are designated with increasing distance away from a paleoshoreline, it cannot be assumed that because an organism exists in zones A and C that it was ever present in zone B. Consequently, the probabilities generated from the methods described below are relative, and cannot be taken as a “true” probability. However, the methods

utilized were designed to be as inclusive as possible and to deliver a strongly conservative estimate of true sampling probabilities between groups; consequently, these methods provide a useful estimate on the relative likelihood of sampling aragonitic or calcitic fauna. Furthermore, sampling probabilities through time based on regional studies such as those utilized by Foote et al. (2015) rely on the assumption that groups were not genuinely absent from the study region at a particular time and that other geographic variables do not have an effect – as such the use of these metrics to evaluate the distribution of fauna across the WIS is validated.

Three methods were devised for dealing with the issue of unknown “correct” distribution of species across the seaway and to correct for differences in the number of collections between zones: (1) finds two bins either side of the current bin and generates a list of the total number of possible species across those five bins; (2) finds all formations that appear in the selected bin that contain specimens of the selected group (e.g. calcitic bivalves), and then finds the total number of species for that group from those formations; (3) finds all formations in the current bin and the two adjacent bins that contain specimens of the selected group, and subsequently finds the total number of species from those formations. For all three methods, the total number of sampling opportunities per bin was generated by multiplying the number of potentially recoverable species by the number of collections to standardize for differences in collecting intensity. The low number of depth-based bins could potentially result in flattening the curve of sampling probability using Method 3, and thus Method 2 is employed in the main body of this paper for depth-based results.

To determine the primary controls on sampling probability between the two stages, we used multiple logistic regression, coding sampling opportunities as the response variable and mineralogy, lithology, life habits (mobility, relation to substrate and feeding style) and depth zone as the predictor variables. Multiple logistic regression allows for the use of binomial

nominal values by using the odds ratio, a measure of the relationship between the odds of an outcome, in this case sampled (1) or not sampled (0), along with multiple potentially explanatory ecological or physiographic variables. A full model is generated that incorporates all potential variables, and a null model defined that includes none. Stepwise addition or deletion from the null or full models, respectively, and analysis in the change of likelihood and of respective AIC (Akaike information criterion) scores contributes to a final predictive model of explanatory variables and respective statistical significance.

Sampling opportunities were tabulated as the presence or absence of each recoverable genera per collection, per depth zone. Each sampling opportunity was assigned a lithology based on collection lithology, as well as all ecological attributes related to that genus. To test for multicollinearity between variables, correlation tests were run using Spearman-Rank correlation using the Performance Analytics package in R. Explanatory variables that showed a strong (above 0.7) statistically significant correlation were excluded from further analysis (Supplementary Information 1).

Interaction terms were also added to explore the possibility of multiple confounding factors and increased model complexity. These terms were restricted to a combination of lithology and mineralogy, so as to test for specific interactions between the two (e.g. whether preservation of aragonite was specifically enhanced within limestones). We also partitioned the data to be able to fully explore the influence of various contributing factors on sampling probability per depth zone, as well as include all organisms in the data (ammonites were excluded from analyses involving life habits, as discussed below). Both effect sizes of individual factors and AIC values of models are presented for statistically significant interactions. All methods were written and implemented using R.

Occurrences, Raw diversity and SQS

To establish the potential influence of aragonite bias on diversity of shelly taxa, total occurrences of organisms were counted per zone using the Select By Location tool in ArcGIS (ESRI, 2010) which were used to generate landward-to-basinward profiles of raw occurrences, raw and subsampled diversity estimates. Shareholder quorum subsampling (SQS; Alroy, 2010), a method for standardising taxonomic occurrence lists based on an estimate of coverage, was implemented in R using script provided by Alroy (pers. comms.) for each faunal group. Calcitic and aragonitic groups were evaluated for statistically significant differences using the Chi-squared test for non-random association (Brown and Rothery, 1993). All statistical tests were implemented in R. Results pertaining to patterns within raw occurrences can be found within Supplementary Information 1 and Figure S1.

RESULTS

Sampling Probability

Cenomanian-Turonian

For generic level sampling probability (Fig. 4a), aragonitic bivalves and ammonites show a similar trend for the first three depth zones. After this, sampling probability drops to 0 for aragonitic bivalves (as none were recovered), whilst it increases to a peak for ammonites. Calcitic bivalves record a higher sampling probability than ammonites or aragonitic bivalves in all zones and show a basinwards increase in sampling probability.

Campanian

In the lower Campanian (Fig. 4b) ammonites have the highest sampling probabilities, showing a level trend across the seaway with a pronounced trough in the distal offshore. Aragonitic bivalves record a relative high sampling probability in the nearshore, followed by a sharp decline for both proximal and distal offshore zones and an increase towards the basin center.

Calcitic fauna have a consistently higher sampling probability than aragonitic bivalves, but lower than ammonites; they also show a level trend across the seaway, experiencing a peak in the distal offshore.

Sampling probability between lithologies

Cenomanian-Turonian

For the Cenomanian-Turonian (Fig. 4c,e,g), ammonites show the same trends and relatively little difference in absolute values between carbonate and siliciclastic sampling opportunities; the greatest difference appears in the basin center, where sampling probability is higher in carbonates. Aragonitic bivalves show a much larger difference, with siliciclastic opportunities scoring consistently higher than carbonate opportunities, even during the large decline within the proximal offshore. Calcitic bivalves show virtually no difference in sampling probability until the basin center, where sampling probability within carbonate sampling opportunities increases substantially.

Campanian

For the Campanian (Fig. 4d,f,h), siliciclastic opportunities of ammonites score higher than carbonate except for within the nearshore. Aragonitic bivalves are not sampled within carbonate collections in either the nearshore, distal offshore or basin center; their sampling probability curve is virtually entirely made by appearances in siliciclastic sampling opportunities. Calcitic bivalves show a decoupled trend between lithologies, with carbonate sampling opportunities showing higher on average sampling probabilities that increase towards the basin center, compared to the fairly low scoring, level trend in siliciclastic.

Multiple Logistic Regression

Results of the logistic regressions are shown in Tables 1-8 and summarized in Fig 5. When interpreting these, note that calcitic mineralogy is compared to aragonitic, so that positive regression coefficients indicate greater odds of sampling calcite. As lithology has multiple parameters, these were compared against the relative sampling probability of mudstone, which is used as a baseline. We are primarily interested in reporting effect sizes, which are gauged by the magnitude of regression coefficients.

AIC scores are utilized in choosing ideal model fit when comparing models with and without two-way interactive terms (a combination of effects between explanatory parameters: for example, the relative odds of sampling calcitic fauna within a specific lithology), with lower scores indicating a better model fit. Only models with the lowest AIC scores are presented here and we only report factors with statistically significant results ($p < 0.05$); full results can be found within Supplementary Information 2.

Cenomanian-Turonian

Mineralogy, lithology, feeding style and depth zone all influence the preservation potential of fauna in the seaway (Table 1); lower AIC scores when an interactive term is added suggest this provides a better model fit than when this is excluded. The odds of sampling calcitic fauna are shown to be 4.6 times (the exponential of the coefficient; 1.52) higher than that of aragonitic fauna, with ANOVA results showing mineralogy contributing the most towards deviance from the null model. Limestone environments are shown to be detrimental to the sampling probability of fauna, whereas sandstones and siltstone enhance sampling probability. The positive interaction between mineralogy and limestone lithologies shows that aragonitic fauna have comparatively strongly reduced odds of being sampled within limestone environments. All depth zones are shown to have decreased sampling probability compared to the basin center, with nearshore and proximal offshore zones showing the worst sampling potential.

397 Chemosymbiont deposit feeders are shown to have an increased preservation potential
398 compared to other feeding styles.

399

400 We additionally partitioned the data into each depth zone, to test for differences with increased
401 bathymetry across the seaway (Table 2). The nearshore zone exhibits an increase in the odds
402 of sampling calcitic fauna, although this effect is reduced compared to results across the whole
403 seaway. Sandstones are also shown to exhibit increased sampling probability. The proximal
404 offshore shows a significant increase in the odds of sampling calcitic bivalves relative to
405 aragonitic bivalves (6.17 compared to 1.88 for the nearshore), as well as increased sampling
406 probability in marl depositional environments and for chemosymbiotic deposit feeders.
407 Limestone negatively impacts the sampling probability of bivalves; the positive interaction
408 between calcite and limestone consequently suggests that this negative impact is related to the
409 sampling probability of aragonitic bivalves. The distal offshore shows a similar pattern,
410 although the relative odds of each are reduced compared to the proximal offshore. The basin
411 center shows increased odds of sampling bivalves within calcarenite, calcareous shale and marl
412 environments, but no other statistically significant terms.

413

414 We also assessed depth zones for the inclusion of all organisms (Table 3). When ammonites
415 are included, the odds of sampling aragonitic fauna increase (calcitic bivalves show odds of 2.1
416 higher sampling probability). Sandstone shows reduced odds of sampling any fauna, the
417 opposite of previous results. The interaction between mineralogy and lithology shows
418 increased sampling probability of calcitic organisms within limestones, sandstones,
419 calcarenites and calcareous mudstones, suggesting this effect is predominantly produced by the
420 addition of ammonite fauna.

When assessing zones independently (Table 4), nearshore sampling probabilities are only controlled by mineralogy, although again with lower odds than reported elsewhere (1.56). In the proximal offshore, results show an increased sampling probability of calcitic fauna within sandstones and calcareous mudstones. The distal offshore also shows strong interactions between sampling probability of calcitic fauna and lithology, with strongly positive coefficients for sandstone, limestone, calcareous shale, and calcarenite two-way interactions. Overall, the sampling probability of calcite compared to aragonitic fauna is high, although reduced compared to the proximal offshore. Within the basin center, mineralogy is not listed as a statistically significant interactive term on its own, but calcitic fauna exhibit increased sampling probability for interactive terms with calcarenites, calcareous mudstones, limestones, and marls.

Campanian

Models for all bivalves in the Campanian (Table 5) show comparatively few statistically significant contributors to sampling probability. By comparison with the Cenomanian, bivalve samples from the Campanian are only weakly influenced by mineralogy (showing odds of 2.16 increased likelihood of sampling calcitic organisms). Additionally, only sandstone and interactions between sandstone and limestone with calcitic organisms are shown to exert any other influence on sampling probability.

This trend continues when partitioning the bivalve data into depth zones (Table 6). The nearshore zone has no statistically significant individual factors contributing to sampling probability. The proximal offshore includes statistically significant effects due to mineralogy and lithology, particularly limestones and calcareous mudstones where sampling probability is enhanced. Mineralogy, sandstone and the interaction between mineralogy and sandstone are reported as statistically significant factors for the distal offshore; mineralogy has a relatively

high positive coefficient (odds of 3.16 in favour of calcitic organisms). Sampling probability is enhanced in sandstones overall, but negatively influences the odds of recovering calcitic organisms: it therefore follows that aragonitic bivalves show particularly enhanced sampling within sandstones. Model results for the basin center suggest that only calcareous shale has a statistically significant positive impact on sampling probability.

When all organisms are assessed (Table 7), mineralogy and depth zone are the only contributors to the full model. Surprisingly, aragonitic organisms have a higher sampling probability than calcitic using the full model, with mineralogy only contributing to a very small amount of deviance from the null ANOVA model. As this result is not observed when assessing bivalve fauna, it is likely that ammonite occurrences are principally contributing to this effect. Depth zones were also evaluated for all organisms (Table 8). Only the proximal offshore supported a model other than the null, which reported mineralogy as a contributing factor; unusually, calcitic fauna are shown to have a reduced sampling probability compared to aragonitic.

Range Size

Cenomanian-Turonian

Box plots were generated on a log scale to show differences in mean range sizes between calcitic and aragonitic organisms (Fig. 6a). There is a visible difference in variability of range size between groupings, with calcitic fauna showing an average larger range than aragonitic. The Wilcoxon Mann Whitney test also showed a statistically significant difference between the range sizes for the two groups (p value = 0.00405), with a reported difference in median range size of 48,694 km². As sample size varied between the groups, resampling measures were carried out to test the accuracy of these results. A randomized bootstrap with replacement calculating the difference between the means of range sizes was implemented 10,000 times in

R (Fig. 6c). Our recorded difference in the mean was shown to have an associated p value of 0.0172, showing statistical significance.

Campanian

Box plots were generated to show differences in mean range sizes between early Campanian calcitic and aragonitic organisms (Fig. 6b). Calcitic bivalves show higher variability in mean range size than aragonitic bivalves. However, the Wilcoxon Mann Whitney test showed no statistically significant difference between the two groupings (p value = 0.504) with a recorded difference in median range size of 13,540 km², and a randomized bootstrap (Fig. 6d) with replacement recovered an associated p value of 0.1527 (non-statistically significant).

Raw Diversity and SQS

Cenomanian-Turonian

Within the Cenomanian-Turonian, broadly similar patterns of diversity occur in all groups (Fig. 7a,c) – peak diversity is within the distal offshore, with lowest values in the nearshore and basin center. Calcitic bivalves show proportionally enhanced diversity in the proximal offshore compared to the other faunal groups. These patterns closely align with the number of collections within each zone, but show limited similarity to zoned outcrop area.

Subsampled ammonite and calcitic bivalve diversity show a broadly similar pattern to their raw taxic diversity signals (Fig. 8a,e). The record of aragonitic bivalves (Fig. 8c) is too poor to resolve subsampled diversity for the basin center; however, a slight decline in subsampled generic richness exists in the proximal offshore.

Campanian

Calcitic bivalves and ammonites exhibit a similar pattern in diversity (Fig. 7b,d) although the latter show an increase in the proximal offshore. Aragonitic bivalve diversity has a similar peak in the proximal offshore but declines towards the basin center. None of these trends show similarity to the distribution of collections or outcrop area throughout the seaway. When subsampled (Fig. 8b,d,f), calcitic and aragonitic bivalves are most diverse within the proximal offshore, falling to relative lows within the distal offshore and basin center. Ammonites are most diverse in the nearshore, followed by a decline to a flat profile.

DISCUSSION

Sampling probability and multiple logistic regression

Our results from estimations of sampling probability and subsequent multiple logistic regression suggest that aragonite bias may be present within distinct depth zones of the seaway during the Cenomanian-Turonian. Mineralogy has a strong and statistically significant impact on sampling probability within the proximal and distal offshore bathymetric zones, and shows the highest contribution to deviance from the null model. This is further supported by the fact that whilst all aragonitic taxa have lower sampling proportions overall, both aragonitic bivalves and ammonites disproportionally decrease in sampling probability within the proximal offshore compared to calcitic bivalves. Ammonites, whilst still showing reduced sampling probability compared to calcitic fauna, are more likely to be sampled than aragonitic bivalves; a potential explanation for this difference could be that aragonite dissolution acts differently upon ammonites compared to bivalves. Body sizes of ammonites and bivalves differ, with ammonites generally having larger forms (Jablonski, 1996). This has been known to influence preservation potential and the extent of aragonite dissolution: Wright et al. (2003) showed that ammonites are affected less severely than aragonitic bivalves by early stage aragonite

dissolution, often exhibiting poor preservation rather than complete removal. Our results have the potential to be partially related to this effect.

Aragonitic bivalves have lower absolute sampling probabilities in carbonate environments than in siliciclastic environments, supporting the results of Foote et al. (2015). However, when examining the proximal offshore zone, we can see that sampling probability within siliciclastic lithologies falls dramatically. As this zone records the largest difference in odds of sampling between calcitic and aragonitic taxa, it can be argued that aragonite bias can influence fauna within siliciclastic deposits in epicontinental seas, in contradiction to Foote et al. (2015). The absolute sampling proportions of calcitic bivalves remain relatively consistent (at about 2% of genera per collection) throughout the seaway until the basin center, where they increase dramatically within carbonates compared to siliciclastics. Foote et al. (2015) reported that calcitic organisms experienced higher sampling probabilities in carbonate-rich intervals, which is especially enhanced in limestones. As carbonates make up 93% of total sampling opportunities within this zone, our results align fairly closely with previous findings. Whilst Foote et al. (2015) singled out lithology as an important factor for aragonite dissolution, they did not investigate whether differences in grain size significantly influenced results. Within this study, sandstone and siltstone are consistently shown to have better odds at preserving aragonitic fauna than mudstone. This is unsurprising, considering that coarser, oxidized sediments are likely to contain lower quantities of organic matter than finer sediments, and thus provide less material for the microbial decay which ultimately controls the dissolution of aragonite within the taphonomically active zone (Cherns et al., 2008). However, siltstone appears to have higher odds than sandstone, potentially a reflection of increased quality of preservation in lower energy settings. It should be noted however that only a few models include both siltstone and sandstone and therefore allow for comparison of sampling probabilities.

Potential ecological signals can also be parsed from the results reported here. Within the Cenomanian-Turonian dataset, odds of sampling chemosymbiont deposit feeders within the proximal offshore were higher than for other bivalves, forming a statistically significant part of the final model and accounting for the second highest deviance from the null model. Chemosymbiosis in bivalves occurs in a range of environments to cope with life in sulphide-rich environments, typically at deep sea vents or in sediments at the oxic/anoxic interface (Cavanaugh, 1994). Combined with evidence for poor sampling probability of aragonitic fauna in siliciclastic lithologies, this lends credence to the likelihood of fluctuating benthic oxygen conditions within the proximal offshore, ideal for preferential aragonite dissolution. More broadly, several previous works have suggested that aragonite bias strongly influences perceived trophic communities within molluscan fauna, favouring preservation of specific life habits (Cherns et al., 2008; Cherns and Wright, 2009). Unfortunately, very few statistically significant life habit factors contribute to our final models (Fig. 5), and thus we cannot draw any conclusions regarding preservational shifts in trophic structure. In the basin center, ammonites are more likely to be sampled compared to other organisms. This confirms expectations of enhanced preservation within a predominantly anoxic water column, where dissolution and predation have reduced impact on the removal of fauna emplaced by pelagic fallout (Jordan et al., 2015).

Within the Campanian, there is a somewhat contradictory pattern. Multiple logistic regression results show that mineralogy only has a strong, statistically significant impact on relative sampling odds when assessing bivalves within the proximal and distal offshore bathymetric zones, with only the latter showing a strong deviation from the null model in ANOVA results. When ammonites are added, the odds of sampling aragonitic fauna are actually higher than that of calcitic organisms within the proximal offshore, and all other zones show no statistically significant contributions from mineralogy. This is reinforced when one considers the absolute

proportions of mineralogies sampled: ammonites exhibit the highest overall sampling probability between fauna. A potential cause of this contradiction is preferential sampling bias. Ease of collecting and human interest can result in skewed sampling effort and intensity, potentially inflating (Foote and Sepkoski, 1999) or reducing (Lloyd and Friedman, 2013) the published records of certain taxa, locations, and time periods above others. The WIS has long been known for its abundance and diversity of ammonite fauna, and consequently ammonites have been used for systematic biostratigraphic correlation since the 1930s (Stephenson and Reeside Jr., 1938). An intensive effort to collect ammonites for stratigraphic purposes was carried out by a selection of workers through the latter half of the 20th century to the present day (Scott and Cobban, 1959; Gill and Cobban, 1973; Cobban and Hook, 1984; Cobban et al., 2006; Merewether et al., 2011). Consequently, it is likely that records for biostratigraphically important organisms have been over-inflated compared to other molluscs and between localities. Koch (1978) showed by comparing previously existing collections and newly collected records for the upper Cenomanian *Sciponoceras gracile* zone (now the *Vascoceras diartianum* and *Euomphaloceras septemseriatum* zones; Cobban et al, 2006) that ammonites were better studied and more commonly reported than bivalve fauna. Parts of these collections have made up the majority of the publicly available records of fossil occurrences within the Western Interior, which are utilized in this study. As such, it is possible that ammonites are over-represented in the early Campanian dataset and are skewing perceived results. However, it is still possible to suggest that a suppressed expression of spatial aragonite bias occurs in the distal offshore, albeit at reduced levels in comparison to the Cenomanian-Turonian interval.

Range size

Range size results reported a difference between calcitic and aragonitic bivalves across the two time intervals studied, with aragonitic fauna showing a significantly smaller range size during

the Cenomanian-Turonian but not the Campanian. This variation is also expressed spatially (Fig. 9). Within the Cenomanian-Turonian time slice, aragonitic geographic ranges (Fig. 9a) are generally restricted to the western and northern edges of the seaway in comparison to calcitic geographic ranges, which extend further to the center of the basin, as well as the east and south (Fig. 9c). This same pattern is slightly different in the early Campanian interval (Fig. 9b,d); whilst aragonitic fauna still show a limited range, the difference between both bivalve groups is less pronounced. This pattern also matches with the distribution of carbonate deposition within the WIS: the Cenomanian-Turonian interval experienced widespread carbonate sedimentation – in the form of the Greenhorn Limestone Formation – in the basin center (Miall et al., 2008), whilst deposition in the basin center transitioned from limestones of the Niobrara Formation to the siliciclastic mudstones of the Pierre Shale in the early Campanian (McGookey et al., 1972; Da Gama et al., 2014). As our results confirm that carbonate environments can exacerbate the effects of aragonite dissolution, it is possible that the differences between the Cenomanian-Turonian and the Campanian are partially driven by the enhanced effects of aragonite bias in carbonate-rich environments, resulting in a lowered sampling probability within carbonate-dominated localities.

Occurrence and Diversity Results

Overall, there is some evidence of aragonite dissolution influencing patterns of pure occurrences, taxonomic and subsampled diversity for aragonitic fauna, as previously hypothesized. In the Cenomanian-Turonian, aragonite bias is most pronounced within the proximal offshore bathymetric zone, with a lesser impact within the distal offshore zone. Whilst all fauna show a close correlation to collection counts for depth zones, both aragonitic and calcitic fauna deviate from this correlation in the proximal offshore zone, recording lower raw occurrences and diversity. The same is broadly observed in the Campanian: maximum

disparity of sampling probability between calcitic and aragonitic fauna is observed within the distal offshore zone, where aragonitic occurrences and raw taxic diversity show a noticeable decline and subsequent deviation from sampling proxies. Foote et al. (2015) reported similar results when comparing sampling-corrected results to ones that previously displayed the proportion of aragonitic taxa (Crampton et al., 2006), and concluded that similarities existed between sampling probabilities and relative proportions of aragonitic species.

Despite the potential relationships discussed above, we cannot report conclusive evidence for aragonite bias influencing the sampled diversity of molluscan fauna within the WIS. This aligns with other recent studies showing that despite evidence of widespread aragonite dissolution during early shallow diagenesis, perceived diversity is not largely affected by these processes (Behrensmeyer et al., 2005; Kidwell, 2005; Crampton et al., 2006; Hsieh et al., 2019). Hence, we must additionally look at external influences which might capture, enhance, or control the distribution of aragonitic faunas that would otherwise be lost to preferential dissolution.

Known human influences have potentially contributed to the suppression of aragonite bias on a spatial scale. Whilst the extent to which aragonite dissolution may have influenced our perceived record of molluscan diversity within the WIS is unclear, it is apparent that these records closely correlate with established sampling proxies. Results of Spearmans-rank correlation tests of occurrences and raw taxic diversity against sampling proxies for distance-from-paleoshoreline zones (Table 9) correlate strongly and significantly. It is clear that broader scale sampling trends related to collector effort strongly influence the pattern of faunal distribution across the seaway, potentially overwriting the effects of aragonite dissolution.

Whilst there have been many cases of preferential aragonite dissolution within local studies, aragonitic molluscan fauna are relatively well represented in the global fossil record (Harper, 1998). This paradox suggests that processes must occur which capture records of molluscan fauna at a higher frequency than they are capable of being destroyed. Cherns et al. (2008, 2011)

644 describe “Taphonomic Windows” as events in the fossil record which capture an unbiased view
645 of aragonitic faunas which have escaped preferential dissolution, and detail numerous
646 examples that may have operated within the WIS. One such window that is prevalent within
647 the WIS are concretions, sedimentary mineral masses of varying chemical composition that
648 often form at shallow burial depths early in diagenesis when mineral cement precipitates
649 locally during lithification (Berner, 1968; McCoy et al., 2015). These have the potential to
650 preserve three-dimensional fossilized remains, often in exquisite detail (Dean et al., 2015; Korn
651 and Pagnac, 2017). Concretions are also a characteristic mode of molluscan occurrences within
652 the WIS, with fossil-bearing concretions found commonly throughout the seaway (Landman
653 and Klofak, 2012); as such, they could further contribute to a potential anthropogenic bias in
654 that they provide easily spotted locations to find fauna in otherwise barren strata (such as the
655 Pierre Shale), skewing collection intensity between localities with concretions and those
656 without. However, only ~3% of USGS collections were obtained by selective collecting (Koch,
657 1980), and as USGS records make up ~55% of our finished dataset this suggests that sampling
658 intensity bias might be partially mitigated. Sediment accumulation rate could exert a large
659 influence on the potential for preferential aragonite dissolution to affect spatial zones of the sea
660 floor. If sediment accumulation rates were low, fauna would remain within the TAZ for an
661 extended period of time, and thus are more likely to be removed through physical reworking,
662 bioerosion and enhanced dissolution (Cherns et al., 2011). In contrast, if sediment
663 accumulation rates were high, fauna are likely to have been rapidly buried and thus have
664 escaped into the sub-TAZ region, where vulnerable bioclasts are likely to be stabilized by
665 shallow burial diagenesis (Melim et al., 2002, 2004). Sediment accumulation rates within the
666 WIS varied both longitudinally within a stratigraphic interval (with higher sediment
667 accumulation rates towards the western paleoshoreline) and with increased bathymetry in a
668 single location (Arthur and Sageman, 2005): accounting for this potential influence is

problematic, and the extent of its effects is ambiguous. The result of these factors is a potential suppression of the spatial influence of aragonite dissolution bias on recorded faunal diversity within the WIS.

Spatial Scale and Influence of Bias

The issue of scale is key to understanding the spatial impact of aragonite dissolution (Kosnik et al., 2011). Foote et al. (2015) recorded preferential aragonite bias within carbonate-rich environments on the regional spatial ($\sim 10^6$ km²) and stage-level temporal (1-10 Myr) scales. However, others (Behrensmeyer et al, 2005; Kidwell, 2005; Kiessling et al., 2008; Kosnik et al, 2011) using global-scale data have reported negligible influence of shell mineralogy on temporal trends or frequency of occurrences. Foote et al. (2015) reported three key differences between previous studies and their work: higher taxonomic level of occurrences, larger time bins, and the use of global data. These factors were inferred to “even out” spatial and temporal variations in sampling, mitigating the influence and effect of locally variable biases inherent to the fossil record. Foote et al. (2015) further suggested that as their taxonomic and temporal scales were consistent with previously published work, an increase in spatial scale may prove the most influential factor on demoting the influence of aragonite dissolution.

This result can be easily translated into the spatial expression of aragonite bias by comparing its potential on alpha (within-site), beta (between-site) and gamma (global) diversity. At the alpha level, the impact of aragonite bias on a single species will be at its most severe, particularly within single-bed assemblages (Wright et al, 2003; Bush and Bambach, 2004; Cherns et al 2008, 2011). However, at gamma levels of diversity, the probability of not recording an individual drops substantially due to the number of possible localities to sample from, where various taphonomic windows may result in aragonite preservation. As such, an increased number of localities in a spatial setting are likely to partially obscure localized

694 aragonite dissolution. As we recorded an impact on zoned sampling probabilities and range
695 size of aragonitic fauna in the WIS, but could not conclusively prove an influence on total
696 diversity estimates, our data support the suggestions of Foote et al. (2015) that spatial scale is
697 a dominant factor on the severity of aragonite bias.

698 Whilst unlikely to influence diversity on a global scale, this study has shown that preferential
699 aragonite dissolution has the capacity to govern the sampling probability of a species in
700 geographic space, and thus can influence the ‘variation’ definition of beta diversity (Anderson
701 et al., 2011). As the preferential dissolution of aragonite is a process that is exacerbated by
702 certain environments (Foote et al. 2015), its influence will impact localities with different
703 environmental conditions to differing extents – a species will be lost at one site and recorded
704 at another. Our results confirm this, showing aragonite bias has an effect on observed diversity
705 between locations, at least during times of widespread carbonate deposition.

706 Consequently, when looking at the spatial signal of aragonite dissolution as a whole, we can
707 see a sliding scale of influence: strong, environmentally dependent impact on alpha diversity;
708 a potentially large influence on beta diversity; and a negligible impact on gamma diversity.

709 Bush et al. (2004) grouped biases affecting spatially organized biodiversity in similar alpha,
710 beta and gamma levels, with alpha biases influencing within site diversity and beta and gamma
711 arising from failure to sample all available habitats or environments within a region. Whilst it
712 was noted in this study that taphonomic effects were not included in this definition, this system
713 can be modified in the light of our results. Aragonite bias, whilst operating at an alpha bias
714 (local) level, evidently has the capacity to systematically influence estimates of beta diversity.

715 As such, the influence of some taphonomic biases may be dependent on the spatial scale at
716 which they are observed. This is an important consideration for studies of the spatial
717 distribution of bias in the fossil record (Barnosky et al., 2005; Vilhena and Smith, 2013; Benson
718 et al., 2016; Close et al., 2017), and for paleobiogeographic studies in general.

CONCLUSIONS

- 1) A multifaceted approach shows that preferential aragonite dissolution is spatially variable and impacts on the relative likelihood, absolute sampling probabilities, and range sizes of aragonitic organisms within the Cretaceous Western Interior Seaway of North America for a time interval that straddles the Cenomanian-Turonian boundary. A similar but reduced effect is additionally observed within an early Campanian time interval. A combination of depositional lithology (a limestone dominated basin within the Cenomanian-Turonian compared to a more siliciclastic setting in the early Campanian) and an anoxic basin center are hypothesized as drivers for this effect.
- 2) Carbonate environments enhance the effects of aragonite dissolution and the preservation of calcitic organisms, as has been previously demonstrated. However, in contrast to previous studies, siliciclastic environments are also shown to be influenced by preferential aragonite dissolution.
- 3) Whilst similarities are observed between faunal distribution and absolute sampling probabilities, we cannot conclusively say that aragonite dissolution has influenced perceived diversity of molluscs within the Western Interior Seaway. “Taphonomic windows” act to preserve records of organisms that would otherwise be lost. Other anthropogenic and geologic biases appear to have a more obvious effect on the molluscan record, and likely mask the influence of aragonite dissolution.
- 4) Whilst aragonite bias can be thought of as an “alpha bias”, results show it could have a systematic and severe impact on beta diversity. This suggests that taphonomic biases can act differently at different scales in the spatial realm.

Acknowledgements

We would like to thank A. Quallington and Getech Group plc for permission to use and edit the paleogeographic reconstructions in this study. We are especially thankful to S. Kidwell for providing molluscan mineralogy information, J. Alroy for providing the SQS function in R, as well as C. McKinney and K. Hollis for providing valuable museum datasets. We thank W. Kiessling, A. Bush and one anonymous reviewer for comments which improved an earlier version of this work. We would also like to thank P. Wright, M. Sephton, S. Maidment and A. Chiarenza for fruitful discussions regarding the manuscript, as well as contributors to PhyloPic.org and the PBDB. CDD was supported by a NERC studentship (1394514). This is Paleobiology Database contribution number XXX.

References Cited

- Allison, P.A., 1988. The role of anoxia in the decay and mineralization of proteinaceous macrofossils. *Paleobiology*, 14, 139–154.
- Allison, P.A., Bottjer, D.J., 2011. Taphonomy: bias and process through time, in: *Taphonomy*. Springer, pp. 1–17.
- Allison, P.A., Wells, M.R., 2006. Circulation in Large Ancient Epicontinental Seas: What Was Different and Why? *Palaaios*, 21, 513–515. <https://doi.org/10.2110/palo.2006.S06>
- Alroy, J., Marshall, C.R., Bambach, R.K., Bezusko, K., Foote, M., Fursich, F.T., Hansen, T.A., Holland, S.M., Ivany, L.C., Jablonski, D., Jacobs, D.K., Jones, D.C., Kosnik, M.A., Lidgard, S., Low, S., Miller, A.I., Novack-Gottshall, P.M., Olszewski, T.D., Patzkowsky, M.E., Raup, D.M., Roy, K., Sepkoski, J.J., Jr., Sommers, M.G., Wagner, P.J., Webber, A., 2001. Effects of sampling standardization on estimates of Phanerozoic marine diversification. *Proceedings of the National Academy of Sciences of the United States of America*, 98, 6261–6. <https://doi.org/10.1073/pnas.111144698>
- Alroy, J., 2010. Fair sampling of taxonomic richness and unbiased estimation of origination and extinction rates. *Quantitative methods in paleobiology: Paleontological Society Papers*, 16, 55–80.
- Anderson, M.J., Crist, T.O., Chase, J.M., Vellend, M., Inouye, B.D., Freestone, A.L., Sanders, N.J., Cornell, H.V., Comita, L.S., Davies, K.F., 2011. Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist. *Ecology letters*, 14, 19–28.
- Arthur, M.A., Sageman, B.B., 2005. Sea-level control on source-rock development: perspectives from the Holocene Black Sea, the mid-Cretaceous Western Interior Basin of North America, and the Late Devonian Appalachian Basin. in: *The deposition of organic-carbon-rich sediments: models, mechanisms and consequences*. Harris, N. B. (Ed.), SEPM Special Publication 82, pp. 35–59.
- Barnosky, A.D., Carrasco, M.A., Davis, E.B., 2005. The impact of the species–area relationship on estimates of paleodiversity. *PLoS biology*, 3, e266.
- Behrensmeyer, A.K., Fursich, F.T., Gastaldo, R.A., Kidwell, S.M., Kosnik, M.A., Kowalewski, M., Plotnick, R.E., Rogers, R.R., Alroy, J., 2005. Are the most durable shelly taxa also the most common in the marine fossil record? *Paleobiology*, 31, 607–623.

- Benson, R.B., Butler, R.J., Alroy, J., Mannion, P.D., Carrano, M.T., Lloyd, G.T., 2016. Near-stasis in the long-term diversification of Mesozoic tetrapods. *PLoS biology*, 14, e1002359.
- Berner, R.A., 1968. Calcium carbonate concretions formed by the decomposition of organic matter. *Science*, 159, 195–197.
- Best, M.M., 2008. Contrast in preservation of bivalve death assemblages in siliciclastic and carbonate tropical shelf settings. *Palaaios*, 23, 796–809.
- Best, M.M., Kidwell, S.M., 2000. Bivalve taphonomy in tropical mixed siliciclastic-carbonate settings. II. Effect of bivalve life habits and shell types. *Paleobiology*, 26, 103–115.
- Brett, C.E., Baird, G.C., 1986. Comparative taphonomy: a key to paleoenvironmental interpretation based on fossil preservation. *Palaaios*, 207–227.
- Briggs, D.E., 2003. The role of decay and mineralization in the preservation of soft-bodied fossils. *Annual Review of Earth and Planetary Sciences*, 31, 275–301.
- Brown, D., Rothery, P., 1993. Models in biology: mathematics, statistics and computing. John Wiley & Sons Ltd, Chichester, U.K. pp. 688.
- Bush, A.M., Bambach, R.K., 2004. Did Alpha Diversity Increase during the Phanerozoic? Lifting the Veils of Taphonomic, Latitudinal, and Environmental Biases. *The Journal of Geology*, 112, 625–642.
- Bush, A.M., Markey, M.J., Marshall, C.R., 2004. Removing bias from diversity curves: the effects of spatially organized biodiversity on sampling-standardization. *Paleobiology*, 30, 666–686. [https://doi.org/10.1666/0094-8373\(2004\)030<0666:RBFDC>2.0.CO;2](https://doi.org/10.1666/0094-8373(2004)030<0666:RBFDC>2.0.CO;2)
- Canfield, D.E., Raiswell, R., 1991. Carbonate precipitation and dissolution, in: Allison, P.A., Briggs, D.E.G. (Eds.), *Taphonomy: Releasing the Data Locked in the Fossil Record*. Plenum Press, New York, pp. 411–453.
- Carter, J.G., 1990. Evolutionary significance of shell micro-structure in the Palaeotaxodonta, Pteriomorpha and Isofilibranchia (Bivalvia: Mollusca). *Skeletal biomineralization: patterns, processes and evolutionary trends*, 1, 135–296.
- Cavanaugh, C.M., 1994. Microbial Symbiosis: Patterns of Diversity in the Marine Environment. *Integr Comp Biol*, 34, 79–89. <https://doi.org/10.1093/icb/34.1.79>
- Cherns, L., Wheeley, J.R., Wright, V.P., 2011. Taphonomic bias in shelly faunas through time: early aragonitic dissolution and its implications for the fossil record, in: *Taphonomy*. Allison, P.A., Bottjer, D. J. (Eds.). Springer, pp. 79–105.
- Cherns, L., Wheeley, J.R., Wright, V.P., 2008. Taphonomic windows and molluscan preservation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 270, 220–229.
- Cherns, L., Wright, V.P., 2009. Quantifying the impacts of early diagenetic aragonite dissolution on the fossil record. *PALAIOS*, 24(11), 756–771.
- Cherns, L., Wright, V.P., 2000. Missing molluscs as evidence of large-scale, early skeletal aragonite dissolution in a Silurian sea. *Geology*, 28, 791–794.
- Close, R.A., Benson, R.B., Upchurch, P., Butler, R.J., 2017. Controlling for the species-area effect supports constrained long-term Mesozoic terrestrial vertebrate diversification. *Nature Communications*, 8, 15381.
- Cobban, W.A., Hook, S.C., 1984. Mid-Cretaceous molluscan biostratigraphy and paleogeography of southwestern part of Western Interior, United States. *Jurassic-Cretaceous Biochronology and Paleogeography of North America. Geological Association of Canada Special Paper 27*, 257–271.
- Cobban, W.A., Walaszczyk, I., Obradovich, J.D., McKinney, K.C., 2006. A USGS Zonal Table for the Upper Cretaceous Middle Cenomanian–Maastrichtian of the Western Interior of the United States Based on Ammonites, Inoceramids, and Radiometric Ages, U.S. Geological Survey Open-File Report 2006-1250.
- Crampton, J.S., Foote, M., Beu, A.G., Maxwell, P.A., Cooper, R.A., Matcham, I., Marshall, B.A., Jones, C.M., 2006. The ark was full! Constant to declining Cenozoic shallow marine biodiversity on an isolated midlatitude continent. *Paleobiology*, 32, 509–532. <https://doi.org/10.1666/06014.1>

- 837 Da Gama, R.O., Lutz, B., Desjardins, P., Thompson, M., Prince, I., Espejo, I., 2014. Integrated
838 paleoenvironmental analysis of the Niobrara Formation: Cretaceous Western Interior Seaway,
839 northern Colorado. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 413, 66–80.
- 840 Davies, D.J., Powell, E.N., Stanton, R.J., 1989. Relative rates of shell dissolution and net sediment
841 accumulation—a commentary: can shell beds form by the gradual accumulation of biogenic
842 debris on the sea floor? *Lethaia*, 22, 207–212.
- 843 Dean, C.D., Sutton, M.D., Siveter, Derek J., Siveter, David J., 2015. A novel respiratory architecture
844 in the Silurian mollusc *Acaenoplax*. *Palaeontology*, 58, 839–847.
- 845 ESRI, 2010. ArcGIS—a complete integrated system. Environmental Systems Research Institute, Inc
846 Redlands, California.
- 847 Foote, M., Crampton, J.S., Beu, A.G., Nelson, C.S., 2015. Aragonite bias, and lack of bias, in the
848 fossil record: lithological, environmental, and ecological controls. *Paleobiology*, 41, 245–265.
- 849 Foote, M., Raup, D.M., 1996. Fossil Preservation and the Stratigraphic Ranges of Taxa. *Paleobiology*,
850 121–140.
- 851 Foote, M., Sepkoski, J.J., 1999. Absolute measures of the completeness of the fossil record. *Nature*,
852 398, 415–417.
- 853 Gill, J.R., Cobban, W.A., 1973. Stratigraphy and geologic history of the Montana Group and
854 equivalent rocks, Montana, Wyoming, and North and South Dakota. U.S. Government
855 Printing Office. pp. 37.
- 856 Harper, E.M., 2000. Are calcitic layers an effective adaptation against shell dissolution in the
857 Bivalvia? *Journal of Zoology*, 251, 179–186. [https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-7998.2000.tb00602.x)
858 [7998.2000.tb00602.x](https://doi.org/10.1111/j.1469-7998.2000.tb00602.x)
- 859 Harper, E.M., 1998. The fossil record of bivalve molluscs. The adequacy of the fossil record. Wiley,
860 Chichester, UK, pp. 243–267.
- 861 Hendy, A.J., 2011. Taphonomic overprints on Phanerozoic trends in biodiversity: Lithification and
862 other secular megabiases, in: *Taphonomy*. Springer, pp. 19–77.
- 863 Hollis, K.A., 2008. Using taphonomic disparity to understand preservation biases in the Western
864 Interior Seaway: An example from the Pierre Shale (Upper Cretaceous). University of
865 Colorado at Boulder. (MSc). University of Colorado. pp. 63.
- 866 Hsieh, S., Bush, A.M., Bennington, J.B., 2019. Were bivalves ecologically dominant over
867 brachiopods in the late Paleozoic? A test using exceptionally preserved fossil assemblages.
868 *Paleobiology*, 45(2), 265–279.
- 869 Jordan, N., Allison, P.A., Hill, J., Sutton, M.D., 2015. Not all aragonitic molluscs are missing:
870 taphonomy and significance of a unique shelly lagerstätte from the Jurassic of SW Britain.
871 *Lethaia*, 48, 540–548.
- 872 Kidwell, S.M., 2005. Shell composition has no net impact on large-scale evolutionary patterns in
873 mollusks. *Science*, 307, 914–7. <https://doi.org/10.1126/science.1106654>
- 874 Kidwell, S.M., Bosence, D.W., 1991. Taphonomy and time-averaging of marine shelly faunas, in:
875 Allison, P.A., Briggs, D.E.G. (Eds.), *Taphonomy: Releasing the Data Locked in the Fossil*
876 *Record*. Plenum Press, New York, pp. 115–209.
- 877 Kidwell, S.M., Brenchley, P.J., 1994. Patterns in bioclastic accumulation through the Phanerozoic:
878 Changes in input or in destruction? *Geology*, 22, 1139–1143.
- 879 Kidwell, S.M., Jablonski, D., 1983. Taphonomic Feedback Ecological Consequences of Shell
880 Accumulation, in: *Biotic Interactions in Recent and Fossil Benthic Communities*, Topics in
881 *Geobiology*. Springer, Boston, MA, pp. 195–248. [https://doi.org/10.1007/978-1-4757-0740-](https://doi.org/10.1007/978-1-4757-0740-3_5)
882 [3_5](https://doi.org/10.1007/978-1-4757-0740-3_5)
- 883 Kiessling, W., Aberhan, M., Villier, L., 2008. Phanerozoic trends in skeletal mineralogy driven by
884 mass extinctions. *Nature Geoscience*, 1, 527.
- 885 Koch, C.F., 1980. Bivalve species duration, areal extent and population size in a Cretaceous sea.
886 *Paleobiology*, 6, 184–192.
- 887 Koch, C.F., 1978. Bias in the Published Fossil Record. *Paleobiology*, 4, 367–372.
- 888 Korn, A.S., Pagnac, D.C., 2017. Anatomically Linked Occurrence of Concretionary Preservation of
889 Mosasaurs from the Pierre Shale of South Dakota. *Palaaios*, 32, 330–336.

- Kosnik, M.A., Alroy, J., Behrensmeier, A.K., Fürsich, F.T., Gastaldo, R.A., Kidwell, S.M., Kowalewski, M., Plotnick, R.E., Rogers, R.R., Wagner, P.J., 2011. Changes in shell durability of common marine taxa through the Phanerozoic: evidence for biological rather than taphonomic drivers. *Paleobiology*, 37, 303–331. <https://doi.org/10.1666/10022.1>
- Ku, T., Walter, L., Coleman, M., Blake, R., Martini, A., 1999. Coupling between sulfur recycling and syndepositional carbonate dissolution: evidence from oxygen and sulfur isotope composition of pore water sulfate, South Florida Platform, USA. *Geochimica et Cosmochimica Acta*, 63, 2529–2546.
- Landman, N.H., Klofak, S.M., 2012. Anatomy of a concretion: life, death, and burial in the Western Interior Seaway. *Palaaios*, 27, 672–693.
- Lloyd, G.T., Friedman, M., 2013. A survey of palaeontological sampling biases in fishes based on the Phanerozoic record of Great Britain. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 372, 5–17. <https://doi.org/10.1016/j.palaeo.2012.07.023>
- Lockwood, R., 2003. Abundance not linked to survival across the end-Cretaceous mass extinction: patterns in North American bivalves. *Proc. Natl. Acad. Sci.* 100, 2478–2482.
- Markwick, P.J., 2007. The palaeogeographic and palaeoclimatic significance of climate proxies for data- model comparisons. *Deep-Time Perspect. Clim. Change Marrying Signal Comput. Models Biol. Proxies*, 251–312.
- Majewske, O.P., 1974. Recognition of invertebrate fossil fragments in rocks and thin sections. (PhD), Brill Archive, University of California, pp. 210.
- McCoy, V.E., Young, R.T., Briggs, D.E., 2015. Factors controlling exceptional preservation in concretions. *Palaaios*, 30(4), 272–280
- McGookey, D.P., Haun, J.D., Hale, L.A., Goodell, H.G., McCubbin, D.G., Weimer, R., Wulf, G.R., 1972. Cretaceous system. *Geologic Atlas of the Rocky Mountain Region: Rocky Mountain Association of Geologists*, pp. 190–228.
- Melim, L.A., Swart, P.K., Eberli, G.P., 2004. Mixing-zone diagenesis in the subsurface of Florida and the Bahamas. *Journal of Sedimentary Research*, 74, 904–913.
- Melim, L.A., Westphal, H., Swart, P.K., Eberli, G.P., Munnecke, A., 2002. Questioning carbonate diagenetic paradigms: evidence from the Neogene of the Bahamas. *Marine Geology*, 185, 27–53.
- Merewether, E.A., Cobban, W.A., Obradovich, J.D., 2011. Biostratigraphic Data from Upper Cretaceous Formations--eastern Wyoming, Central Colorado, and Northeastern New Mexico. US Department of the Interior, US Geological Survey. USGS Scientific Investigation Map, v. 3175, US Department of the Interior, pp.11
- Miall, A.D., Catuneanu, O., Vakarelov, B.K., Post, R., 2008. The Western interior basin. *Sedimentary basins of the world*, 5, 329–362.
- Myers, C.E., Lieberman, B.S., 2011. Sharks that pass in the night: using Geographical Information Systems to investigate competition in the Cretaceous Western Interior Seaway. *Proceedings. Biological sciences / The Royal Society*, 278, 681–9. <https://doi.org/10.1098/rspb.2010.1617>
- Paul, C.R.C., Allison, P.A., Brett, C.E., 2008. The occurrence and preservation of ammonites in the Blue Lias Formation (lower Jurassic) of Devon and Dorset, England and their palaeoecological, sedimentological and diagenetic significance. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 270(3–4), 258–272.
- Peters, S.E., 2009. The problem with the Paleozoic. *Paleobiology*, 33(2), 165–181
- Raup, D.M., 1976. Species diversity in the Phanerozoic: an interpretation. *Paleobiology*, 289–297.
- Ritterbush, K.A., Hoffmann, R., Lukeneder, A., De Baets, K., 2014. Pelagic palaeoecology: the importance of recent constraints on ammonoid palaeobiology and life history. *Journal of Zoology*, 292, 229–241.
- Ros Franch, S., 2009. Dinámica de la paleodiversidad de los Bivalvos del Triásico y Jurásico Inferior. (PhD), Universidad Valencia, pp. 576.
- Sageman, B.B., Arthur, M.A., 1994. Early Turonian paleogeographic/paleobathymetric map, Western Interior, U.S., in: Caputo, M.V., Peterson, J.A., Franczyk, K.J. (Eds.), *Mesozoic Systems of the Rocky Mountain Region, USA*. pp. 458–469.

Sanders, D., 2003. Syndepositional dissolution of calcium carbonate in neritic carbonate environments: geological recognition, processes, potential significance. *Journal of African Earth Sciences*, 36, 99–134.

Schneider, J.A., Carter, J.G., 2001. Evolution and phylogenetic significance of cardioidean shell microstructure (Mollusca, Bivalvia). *J. Paleontol.* 75, 607–643.

Scott, G.R., Cobban, W.A., 1959. So-called Hygiene group of northeastern Colorado. in: The Rocky Mountain Associate Geologists Guidebook. Symposium on Cretaceous rocks of Colorado and adjacent areas. 11th Field conference Washakie, Sand Wash and Pieceance Basins, pp. 124-131.

Stephenson, L.W., Reeside Jr, J.B., 1938. Comparison of Upper Cretaceous deposits of Gulf region and western interior region. *AAPG Bulletin*, 22, 1629–1638.

Taylor, J.D., 1969. The shell structure and mineralogy of the Bivalvia. Introduction. Nuculacea-Trigonacea. *Bull Br Mus Nat HistZool*, 3, 1–125.

Taylor, J.D., Layman, M., 1972. The mechanical properties of bivalve (Mollusca) shell structures. *Palaeontology*, 15.

Team, R.C., 2014. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2014.

Todd, J.A., 2017. NMITA Molluscan Life Habits databases [WWW Document]. URL <http://porites.geology.uiowa.edu/database/mollusc/mollusclifestyles.htm> (accessed 1.11.18).

Tynan, S., Opdyke, B.N., 2011. Effects of lower surface ocean pH upon the stability of shallow water carbonate sediments. *Science of the Total Environment*, 409, 1082–1086.

Valentine, J.W., Jablonski, D., Kidwell, S., Roy, K., 2006. Assessing the fidelity of the fossil record by using marine bivalves. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 6599–604. <https://doi.org/10.1073/pnas.0601264103>

Vilhena, D.A., Smith, A.B., 2013. Spatial Bias in the Marine Fossil Record. *PLoS ONE*, 8, e74470. <https://doi.org/10.1371/journal.pone.0074470>

Walter, L.M., Bischof, S.A., Patterson, W.P., Lyons, T.W., O’Nions, R., Gruszczynski, M., Sellwood, B., Coleman, M., 1993. Dissolution and recrystallization in modern shelf carbonates: evidence from pore water and solid phase chemistry [and discussion]. *Philosophical Transactions of the Royal Society of London. Series A: Physical and Engineering Sciences*, 344, 27–36.

Walter, L.M., Morse, J.W., 1984. Reactive surface area of skeletal carbonates during dissolution; effect of grain size. *Journal of Sedimentary Research*, 54, 1081–1090. <https://doi.org/10.1306/212F8562-2B24-11D7-8648000102C1865D>

Wright, P., Cherns, L., Hodges, P., 2003. Missing molluscs: Field testing taphonomic loss in the Mesozoic through early large-scale aragonite dissolution. *Geology*, 31, 211. [https://doi.org/10.1130/0091-7613\(2003\)031<0211:mmfttl>2.0.co;2](https://doi.org/10.1130/0091-7613(2003)031<0211:mmfttl>2.0.co;2)

Figure and Table Captions

Fig. 1.

Diagram showing potential model of spatial aragonite bias within the WIS. Within the outer shelf, preferential dissolution of aragonitic fauna is common, which has the potential to be expressed spatially. Within the basin center, anoxia limits benthic organism development, but allows for preservation of aragonitic material. Modified after Cherns et al. (2011).

Fig. 2.

Paleogeographic zoned maps of the WIS used in this study. Depth-based zones are designated as nearshore, proximal offshore, distal offshore and basin center (Fig. 1); A. Paleobathymetric map of the Cenomanian-Turonian; B. Paleobathymetric map of the early Campanian.

Fig. 3.

Diagram showing potential model of apparent range size reduction due to spatially variable aragonite preservation. Assuming that calcitic and aragonitic species of bivalve were both living at four separate localities but aragonitic dissolution strongly influenced one of those locations (A), the resulting convex hull for the aragonitic fauna drawn from surviving fossil occurrences would likely be smaller than that of the calcitic organism (B).

Fig. 4.

Plots of generic level M2 sampling probabilities for the Cenomanian-Turonian (A, C, E, G) and lower Campanian (B, D, F, H) time slices across depth zones, split into carbonate and siliciclastic sampling opportunities. All results are plotted with percentage of carbonate collections per depth zone A. Cenomanian-Turonian generic level sampling probability, plotted with percentage of carbonate collections per depth zone; B. lower Campanian generic level sampling probability, plotted with percentage of carbonate collections per depth zone; C. Cenomanian-Turonian ammonite sampling probability; D. lower Campanian ammonite sampling probability; E. Cenomanian-Turonian aragonitic bivalve sampling probability; F. lower Campanian aragonitic bivalve sampling probability; E. Cenomanian-Turonian calcitic bivalve sampling probability; F. lower Campanian calcitic bivalve sampling probability.

Fig. 5

Graph summarizing multiple logistic regression model results (T1-T8 = Tables 1-8). Final models are presented within columns, whereas factors are presented along rows for those models. Results are presented as either green (comparatively positive odds of sampling compared to reference factor), red (comparatively negative odds of sampling compared to reference factor), light grey (factor not statistically significant) and dark grey (factor not included in the final model). The following factors are used as a baseline for comparison: Mineralogy = aragonite; lithology = mudstone, lithology with mineralogy = aragonite:mudstone; zone = Basin Center (BC); diet = carnivore. Note that the magnitude of regression coefficients is not presented within this graph. NS: Near Shore; POS: Proximal Offshore; DOS: Distal offshore; BC: Basin Center.

Fig. 6.

Range size plots for the Cenomanian-Turonian and lower Campanian. A. Cenomanian-Turonian box plots of range size for both aragonitic bivalves and calcitic bivalves on log scale; B. lower Campanian box plots of range size for both aragonitic bivalves and calcitic bivalves on log scale; C. Randomized bootstrap for Cenomanian-Turonian mean range sizes – recorded difference in the mean is shown to be statistically significant; D. Randomized bootstrap for lower Campanian mean range sizes – recorded difference in the mean is not shown to be statistically significant.

Fig. 7.

Plots of generic level diversity plots for the Cenomanian-Turonian and lower Campanian within depth zones, plotted with number of collections and outcrop area. A. Generic diversity and number of collections for the Cenomanian-Turonian; B. Generic diversity and number of collections for the lower Campanian; C. Generic diversity and outcrop area for the Cenomanian-Turonian; D. Generic diversity and outcrop area for the lower Campanian.

Fig. 8.

Plots of generic level SQS results for depth zones in the Cenomanian-Turonian and lower Campanian, set at 0.4, 0.5 and 0.6 quora. A. SQS results for ammonites in the Cenomanian-Turonian; B. SQS

results for ammonites in the lower Campanian; C. SQS results for aragonitic bivalves in the Cenomanian-Turonian; D. SQS results for aragonitic bivalves in the lower Campanian; E. SQS results for calcitic bivalves in the Cenomanian-Turonian; F. SQS results for calcitic bivalves in the lower Campanian.

Fig. 9.

Paleogeographic maps shown with range sizes of calcitic and aragonitic bivalves for both time slices. A. Aragonitic bivalve range sizes for the Cenomanian-Turonian; B. Aragonitic bivalve range sizes for the lower Campanian; C. Calcitic bivalve range sizes for the Cenomanian-Turonian; D. Calcitic bivalve range sizes for the lower Campanian.

Table 1.

Table for Multiple Logistic Regression results for all bivalves within the Cenomanian-Turonian across the whole seaway, using model with lowest AIC score. M - Mineralogy; D - Diet; L - Lithology; Z – depth zone.

Table 2.

Table for Multiple Logistic Regression results for all bivalves within the Cenomanian-Turonian for each depth zone, using models with lowest AIC scores. M - Mineralogy; D - Diet; L – Lithology.

Table 3.

Table for Multiple Logistic Regression results for all organisms (including ammonites) within the Cenomanian-Turonian across the whole seaway, using model with lowest AIC score. M - Mineralogy; L - Lithology; Z – depth zone.

Table 4.

Table for Multiple Logistic Regression results for all organisms (including ammonites) within the Cenomanian-Turonian for each depth zone, using models with lowest AIC scores. M - Mineralogy; L – Lithology.

Table 5.

Table for Multiple Logistic Regression results for all bivalves within the lower Campanian across the whole seaway, using model with lowest AIC score. M - Mineralogy; D - Diet; L - Lithology; Z – depth zone.

Table 6.

Table for Multiple Logistic Regression results for all bivalves within the lower Campanian for each depth zone, using models with lowest AIC scores. M - Mineralogy; D - Diet; L - Lithology; Z – depth zone.

Table 7.

Table for Multiple Logistic Regression results for all organisms (including ammonites) within the lower Campanian across the whole seaway, using model with lowest AIC score. M - Mineralogy; L - Lithology; Z – depth zone.

Table 8.

1104 Table for Multiple Logistic Regression results for all organisms (including ammonites) within the lower
1105 Campanian for each depth zone, using models with lowest AIC scores. M - Mineralogy; L – Lithology.
1106
1107 Table 9.
1108
1109 Spearman's rank correlations between generic diversity of faunal groups and various sampling proxies
1110 for distance-from-paleoshoreline zones within the Cenomanian-Turonian and lower Campanian.

For Peer Review




Figure 1. Diagram showing potential model of spatial aragonite bias within the WIS. Within the outer shelf, preferential dissolution of aragonitic fauna is common, which has the potential to be expressed spatially. Within the basin center, anoxia limits benthic organism development, but allows for preservation of aragonitic material. Modified after Cherns et al. (2011).




Figure 2. Paleogeographic zoned maps of the WIS used in this study. Depth-based zones are designated as nearshore, proximal offshore, distal offshore and basin center (Fig. 1); A. Paleobathymetric map of the Cenomanian-Turonian; B. Paleobathymetric map of the early Campanian.







Figure 3. Diagram showing potential model of apparent range size reduction due to spatially variable aragonite preservation. Assuming that calcitic and aragonitic species of bivalve were both living at four separate localities but aragonitic dissolution strongly influenced one of those locations (A), the resulting convex hull for the aragonitic fauna drawn from surviving fossil occurrences would likely be smaller than that of the calcitic organism (B).




Plots of generic level M2 sampling probabilities for the Cenomanian-Turonian (A, C, E, G) and lower Campanian (B, D, F, H) time slices across depth zones, split into carbonate and siliciclastic sampling opportunities. All results are plotted with percentage of carbonate collections per depth zone. A. Cenomanian-Turonian generic level sampling probability, plotted with percentage of carbonate collections per depth zone; B. lower Campanian generic level sampling probability, plotted with percentage of carbonate collections per depth zone; C. Cenomanian-Turonian ammonite sampling probability; D. lower Campanian ammonite sampling probability; E. Cenomanian-Turonian aragonitic bivalve sampling probability; F. lower Campanian aragonitic bivalve sampling probability; G. Cenomanian-Turonian calcitic bivalve sampling probability; H. lower Campanian calcitic bivalve sampling probability.




Graph summarizing multiple logistic regression model results (T1-T8 = Tables 1-8). Final models are presented within columns, whereas factors are presented along rows for those models. Results are presented as either green (comparatively positive odds of sampling compared to reference factor), red (comparatively negative odds of sampling compared to reference factor), light grey (factor not statistically significant) and dark grey (factor not included in the final model). The following factors are used as a baseline for comparison: Mineralogy = aragonite; lithology = mudstone, lithology with mineralogy = aragonite:mudstone; zone = Basin Center (BC); diet = carnivore. Note that the magnitude of regression coefficients is not presented within this graph. NS: Near Shore; POS: Proximal Offshore; DOS: Distal offshore; BC: Basin Center.




Range size plots for the Cenomanian-Turonian and lower Campanian. A. Cenomanian-Turonian box plots of range size for both aragonitic bivalves and calcitic bivalves on log scale; B. lower Campanian box plots of range size for both aragonitic bivalves and calcitic bivalves on log scale; C. Randomized bootstrap for Cenomanian-Turonian mean range sizes – recorded difference in the mean is shown to be statistically significant; D. Randomized bootstrap for lower Campanian mean range sizes – recorded difference in the mean is not shown to be statistically significant.



Plots of generic level diversity plots for the Cenomanian-Turonian and lower Campanian within depth zones, plotted with number of collections and outcrop area. A. Generic diversity and number of collections for the Cenomanian-Turonian; B. Generic diversity and number of collections for the lower Campanian; C. Generic diversity and outcrop area for the Cenomanian-Turonian; D. Generic diversity and outcrop area for the lower Campanian.



Plots of generic level SQS results for depth zones in the Cenomanian-Turonian and lower Campanian, set at 0.4, 0.5 and 0.6 quora. A. SQS results for ammonites in the Cenomanian-Turonian; B. SQS results for ammonites in the lower Campanian; C. SQS results for aragonitic bivalves in the Cenomanian-Turonian; D. SQS results for aragonitic bivalves in the lower Campanian; E. SQS results for calcitic bivalves in the Cenomanian-Turonian; F. SQS results for calcitic bivalves in the lower Campanian.



Paleogeographic maps shown with range sizes of calcitic and aragonitic bivalves for both time slices. A. Aragonitic bivalve range sizes for the Cenomanian-Turonian; B. Aragonitic bivalve range sizes for the lower Campanian; C. Calcitic bivalve range sizes for the Cenomanian-Turonian; D. Calcitic bivalve range sizes for the lower Campanian.

<i>Final Model</i>	<i>AIC</i>	<i>Factor</i>	<i>Regression coefficient</i>	<i>Standard error</i>
M + D + L + Z + M:L	22367*	M-Calcite	1.516	0.080
		D-Chemosymbiont deposit feeder	1.353	0.239
		L-Sandstone	0.672	0.115
		L-Silt	0.759	0.362
		M-Calcite:L-Limestone	1.338	0.192
		L-Limestone	-1.415	0.184
		DOS	-0.410	0.082
		NS	-0.754	0.114
		POS	-0.684	0.089
		D-Herbivore	-1.870	1.022
		M-Calcite:L-Calcarenite	1.146	0.611

Table for Multiple Logistic Regression results for all bivalves within the Cenomanian-Turonian across the who

<i>p-value</i>
< 2e-16 ***
1.45e-08 ***
5.39e-09 ***
0.0359 *
3.35e-12 ***
1.49e-14 ***
6.70e-07 ***
4.30e-11 ***
1.10e-14 ***
0.0673 .
0.0605 .

ple seaway, using model with lowest AIC score. M - Mineralogy; D - Diet; L - Lithology; Z – depth zone.

For Peer Review

For Peer Review

<i>Data</i>	<i>Final Model</i>	<i>AIC</i>	<i>Factor</i>
Near Shore	M + L	1808.7	M-Calcite L-Sandstone
Proximal Offshore	M + D + L + M:L	7809.2	M-Calcite D-Chemosymbiont deposit feeder L-Marl M-Calcite:L-Limestone L-Limestone M-Calcite:L-Marl
Distal Offshore	M + L + D + M:L	10768	M-Calcite L-Sandstone L-Silt D-Chemosymbiont deposit feeder M-Calcite:L-Limestone L-Limestone
Basin Center	M + L + D	1857.1	L-Calcarenite L-Calcareous Shale L-Marl

Table for Multiple Logistic Regression results for all bivalves within the Cenomanian-Turonian

<i>Regression coefficient</i>	<i>Standard error</i>	<i>p-value</i>
0.626	0.176	0.000372 ***
0.571	0.166	0.000593 ***
1.818	0.134	< 2e-16 ***
1.917	0.449	1.94e-05 ***
3.125	0.539	6.89e-09 ***
1.390	0.438	0.00149 **
-1.358	0.422	0.00131 **
-1.775	0.808	0.02804 *
1.416	0.111	< 2e-16 ***
0.978	0.152	1.18e-10 ***
1.684	0.595	0.004671 **
1.051	0.293	0.000335 ***
1.222	0.226	6.08e-08 ***
-1.248	0.211	3.25e-09 ***
1.777	0.582	0.00225 **
1.826	0.371	8.47e-07 ***
1.864	0.489	0.00014 ***

ian for each depth zone, using models with lowest AIC scores. M - Mineralogy; D - Diet; L – Lithology.

For Peer Review

<i>Final Model</i>	<i>AIC</i>	<i>Factor</i>	<i>Regression coefficient</i>	<i>Standard error</i>
M + Z + L + M:L	46464*	M-Calcite	0.741	0.044
		L-Conglomerate	1.495	0.324
		L-Marl	0.593	0.174
		M-Calcite:L-Calcareenite	1.269	0.319
		M-Calcite:L-Calcareous Shale	2.290	0.280
		M-Calcite:L-Limestone	0.202	0.072
		M-Calcite:L-Sandstone	0.595	0.095
		DOS	-0.530	0.050
		NS	-0.700	0.076
		POS	-0.755	0.056
		L-Calcareenite	-0.959	0.260
		L-Calcareous Shale	-1.561	0.253
		L-Limestone	-0.288	0.050
		L-Sandstone	-0.152	0.067
		M-Calcite:L-Conglomerate	-1.857	1.057

Table for Multiple Logistic Regression for all organisms (including ammonites) within the Cenomanian-Turc

<i>p-value</i>
< 2e-16 ***
3.92e-06 ***
0.000654 ***
7.10e-05 ***
3.03e-16 ***
0.004996 **
4.31e-10 ***
< 2e-16 ***
< 2e-16 ***
< 2e-16 ***
0.000227 ***
6.77e-10 ***
6.82e-09 ***
0.024007 *
0.079016 .

onian across the whole seaway, using model with lowest AIC score. M - Mineralogy; L - Lithology; Z – depth zc

one.

For Peer Review

<i>Data</i>	<i>Final Model</i>	<i>AIC</i>	<i>Factor</i>	<i>Regression coefficient</i>
Near Shore	M	3563.3	M-Calcite	0.443
Proximal Offshore	M + L + M:L	14954*	M-Calcite	0.942
			L-Conglomerate	1.553
			L-Marl	2.046
			M-Calcite:L-Calcareous Shale	1.900
			M-Calcite:L-Sandstone	0.368
			L-Calcareous Shale	-1.765
			L-Sandstone	-0.261
			L-Limestone	-0.203
			M-Calcite:L-Conglomerate	-2.057
Distal Offshore	M + L + M:L	21160*	M-Calcite	0.701
			L-Marl	0.822
			M-Calcite:L-Calcareous Shale	1.639
			M-Calcite:L-Calcareous Shale	1.842
			M-Calcite:L-Limestone	0.287
			M-Calcite:L-Sandstone	0.849
			L-Calcareous Shale	-1.530
			L-Calcareous Shale	-1.651
			L-Limestone	-0.313
			M-Calcite:L-Marl	-1.747
			M-Calcite:L-Silt	1.074
Basin Center	M + L + M:L	6692.3*	L-Marl	-2.270
			M-Calcite:L-Calcareous Shale	3.536
			M-Calcite:L-Calcareous Shale	3.479
			M-Calcite:L-Limestone	0.868
			M-Calcite:L-Marl	4.135

Table for Multiple Logistic Regression results for all organisms (including ammonites) within the Cenomanian

<i>Standard error</i>	<i>p-value</i>
0.114	9.92e-05 ***
0.070	< 2e-16 ***
0.325	1.79e-06 ***
0.367	2.47e-08 ***
0.636	0.00281 **
0.153	0.01635 *
0.578	0.00224 **
0.108	0.01584 *
0.108	0.06071 .
1.059	0.05197 .
0.064	< 2e-16 ***
0.235	0.000475 ***
0.583	0.004940 **
0.657	0.005042 **
0.105	0.006111 **
0.141	1.60e-09 ***
0.502	0.002308 **
0.579	0.004370 **
0.067	3.15e-06 ***
0.750	0.019823 *
0.588	0.067947 .
1.011	0.024660 *
1.167	0.002442 **
0.507	6.53e-12 ***
0.376	0.021101 *
1.123	0.000231 ***

an-Turonian for each depth zone, using models with lowest AIC scores. M - Mineralogy; L – Litho

For Peer Review

ology.

<i>Final Model</i>	<i>AIC</i>	<i>Factor</i>	<i>Regression coefficient</i>	<i>Standard error</i>
M + L + M:L	7063.9*	M-Calcite	0.767	0.115
		L-Sandstone	0.594	0.168
		M-Calcite:L-Limestone	1.292	0.538
		M-Calcite:L-Sandstone	-0.496	0.209
		M-Calcite:L-Siltstone	1.838	1.044

Table for Multiple Logistic Regression results for all bivalves within the lower Campanian across the \

For Peer Review

<i>p-value</i>
2.41e-11 ***
0.00041 ***
0.01625 *
0.01781 *
0.07825 .

whole seaway, using model with lowest AIC score. M - Mineralogy; D - Diet; L - Lithology; Z – depth zone.

For Peer Review

For Peer Review

<i>Data</i>	<i>Final Model</i>	<i>AIC</i>	<i>Factor</i>	<i>Regression coefficient</i>
Near Shore	L	886.47	N/A	N/A
Proximal Offshore	M + L	1603.9	M-Calcite	0.808
			L-Calcareous Shale	0.994
			L-Limestone	0.687
Distal Offshore	M + L + M:L	3420.6*	M-Calcite	1.145
			L-Sandstone	0.765
			M-Calcite:L-Sandstone	-0.709
Basin Center	L + D	1158.2	L-Calcareous Shale	0.812

Table for Multiple Logistic Regression results for all bivalves within the lower Campanian for each d

For Peer Review

<i>Standard error</i>	<i>p-value</i>
N/A	N/A
0.184	1.08e-05 ***
0.467	0.03336 *
0.247	0.00545 **
0.169	1.41e-11 ***
0.260	0.00332 **
0.322	0.02768 *
0.222	0.000259 ***

lepth zone, using models with lowest AIC scores. M - Mineralogy; D - Diet; L - Lithology; Z – depth

For Peer Review

h zone.

For Peer Review

<i>Final Model</i>	<i>AIC</i>	<i>Factor</i>	<i>Regression coefficient</i>	<i>Standard error</i>	<i>p-value</i>
Z + M	16712	DOS	-0.323	0.067	1.61e-06 ***
		NS	-0.203	0.095	0.033311 *
		POS	-0.281	0.078	0.000307 ***
		M-Calcite	-0.146	0.054	0.006599 **

Table for Multiple Logistic Regression results for all organisms (including ammonites) within the lower Car

For Peer Review

mpanian across the whole seaway, using model with lowest AIC score. M - Mineralogy; L - Lithology; Z – dep

For Peer Review

th zone.

For Peer Review

<i>Data</i>	<i>Final Model</i>	<i>AIC</i>	<i>Factor</i>	<i>Regression coefficient</i>	<i>Standard error</i>
Near Shore	Rcoll ~ 1	1747.1	N/A	N/A	N/A
Proximal Offshore	M	3683	M-Calcite	-0.307	0.117
Distal Offshore	Rcoll ~ 1	8319.3	N/A	N/A	N/A
Basin Center	L + M + L:M	1151.9*	N/A	N/A	N/A

Table for Multiple Logistic Regression results for all organisms (including ammonites) within the lo

For Peer Review

<i>p-value</i>
N/A
0.00838 **
N/A
N/A

lower Campanian for each depth zone, using models with lowest AIC scores. M - Mineralogy; L – Lith

For Peer Review

ology.

For Peer Review

DISTANCE			Correlation	Spearman's Rho
CENOMANIAN-TURONIAN	AMMONITES	Outcrop Area	Ammonite occurrences vs. Outcrop Area	0.475
			Ammonite raw diversity vs. Outcrop Area	0.704
		Collections	Ammonite occurrences vs. Collections	0.885
			Ammonite raw diversity vs. Collections	0.732
	ARAG. BIVALVES	Outcrop Area	Aragonitic bivalve occurrences vs. Outcrop Area	0.634
			Aragonitic bivalve raw diversity vs. Outcrop Area	0.640
		Collections	Aragonitic bivalve occurrences vs. Collections	0.568
			Aragonitic bivalve raw diversity vs. Collections	0.502
	CALC. BIVALVES	Outcrop Area	Calcitic bivalve occurrences vs. Outcrop Area	0.534
			Calcitic bivalve raw diversity vs. Outcrop Area	0.678
		Collections	Calcitic bivalve occurrences vs. Collections	0.904
			Calcitic bivalve raw diversity vs. Collections	0.738
LOWER CAMPANIAN	AMMONITES	Outcrop Area	Ammonite occurrences vs. Outcrop Area	0.221
			Ammonite raw diversity vs. Outcrop Area	0.343
		Collections	Ammonite occurrences vs. Collections	0.970
			Ammonite raw diversity vs. Collections	0.727
	ARAG. BIVALVES	Outcrop Area	Aragonitic bivalve occurrences vs. Outcrop Area	0.473
			Aragonitic bivalve raw diversity vs. Outcrop Area	0.404
		Collections	Aragonitic bivalve occurrences vs. Collections	0.822
			Aragonitic bivalve raw diversity vs. Collections	0.759
	CALC. BIVALVES	Outcrop Area	Calcitic bivalve occurrences vs. Outcrop Area	0.056
			Calcitic bivalve raw diversity vs. Outcrop Area	0.322
		Collections	Calcitic bivalve occurrences vs. Collections	0.958
			Calcitic bivalve raw diversity vs. Collections	0.879

Table 9: Spearman's rank correlations between generic diversity of faunal groups and various samp

<i>p-value</i>	<i>Sig.</i>
0.05569	<i>N/A</i>
0.00163	*
0.00000	*
0.00084	*
0.00624	*
0.00569	*
0.01740	*
0.04011	*
0.02918	*
0.00277	*
0.00000	*
0.00071	*
0.49070	<i>N/A</i>
0.27623	<i>N/A</i>
0.00000	*
0.01000	*
0.12070	<i>N/A</i>
0.19248	<i>N/A</i>
0.00104	*
0.00418	*
0.86898	<i>N/A</i>
0.30701	<i>N/A</i>
0.00000	*
0.00017	*

oling proxies for distance-from-paleoshoreline zones within the Cenomanian-Turonian and lower Cam

For Peer Review

spanian